



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
V &= 4 \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi - \phi} \int_{a \sin \phi \sec \theta}^{\frac{1}{2}\pi} \rho^2 \sin \phi d\phi d\theta d\rho = \frac{4}{3} a^3 \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi - \phi} (1 - \sin^3 \phi \sec^3 \theta) \sin \phi d\phi d\theta, \\
&= \frac{4}{3} a^3 \left[\frac{1}{2}\pi - \frac{1}{6} + \frac{3}{16} \int_0^{\frac{1}{2}\pi} \log \tan \frac{\phi}{2} d\phi = \frac{3}{2} \pi a^3 - \frac{5}{4} a^3 + \frac{1}{4} a^3 \int_0^{\frac{1}{2}\pi} \log \tan \frac{\phi}{2} d\phi \right], \\
&= \frac{3}{2} \pi a^3 - \frac{5}{4} a^3 - \frac{1}{4} a^3 \sum_{n=1}^{\infty} \frac{2n-1}{[4n-3]^2 [4n-1]^2} = .386a^3, \text{ nearly.}
\end{aligned}$$

On the integration of $\int_0^{\frac{1}{2}\pi} \log \tan \frac{\phi}{2} d\phi$, see the remarks on Prize Problem, No. 123, Calculus.

157. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

Two equal ellipses are tangent to each other at the vertices of the major axes. If one of them be rolled on the other; find (1) the equation and area of the curve described by the vertex, and (2) by the center.

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa., and the PROPOSER.

Let a and b be the semi-axes of both ellipses; B the center of the fixed ellipse; C , the center of the rolling ellipse; P , its vertex; and D , the point of contact; $e^2 = (a^2 - b^2)/a^2$. Let $BE = x$, $BF = m$, $PE = y$, $CF = n$, and $\angle ABC = \angle PCB = \theta$.

Then, $BC = 2a\sqrt{1 - e^2 \sin^2 \theta}$ = twice the length of the perpendicular from B on the tangent at D .

$$m = 2a \cos \theta \sqrt{1 - e^2 \sin^2 \theta}, \quad n = 2a \sin \theta \sqrt{1 - e^2 \sin^2 \theta},$$

$$x = m - PG = 2a \cos \theta \sqrt{1 - e^2 \sin^2 \theta} - a \cos 2\theta,$$

$$y = n - CG = 2a \sin \theta \sqrt{1 - e^2 \sin^2 \theta} - a \sin 2\theta.$$

$$x^2 + y^2 = r^2 = a^2 + 4a^2(1 - e^2 \sin^2 \theta) - 4a^2 \cos \theta \sqrt{1 - e^2 \sin^2 \theta},$$

the equation of the locus of the vertex.

$$\text{Area} = a^2 \int_0^\pi [5 - 4e^2 \sin^2 \theta - 4 \cos \theta \sqrt{1 - e^2 \sin^2 \theta}] d\theta = \pi a^2 (5 - 2e^2) = \pi (3a^2 + 2b^2).$$

$$BC^2 = r^2 = 4a^2(1 - e^2 \sin^2 \theta), \text{ the equation of the locus of the center.}$$

$$\text{Area} = 4a^2 \int_0^\pi (1 - e^2 \sin^2 \theta) d\theta = 2\pi a^2 (2 - e^2) = 2\pi (a^2 + b^2).$$

Also solved by J. SCHEFFER, and G. W. GREENWOOD.

MECHANICS.

147. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

A particle mass m is attached to one end of a string, the other end of which is fixed. It is projected horizontally with such a velocity that it would rise to a position in which

